

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 1/08, C12N 1/06	A1	(11) International Publication Number: WO 95/28409 (43) International Publication Date: 26 October 1995 (26.10.95)														
(21) International Application Number: PCT/US95/04586 (22) International Filing Date: 13 April 1995 (13.04.95) (30) Priority Data: <table border="0"><tr><td>08/227,514</td><td>14 April 1994 (14.04.94)</td><td>US</td></tr><tr><td>08/227,516</td><td>14 April 1994 (14.04.94)</td><td>US</td></tr><tr><td>08/236,536</td><td>2 May 1994 (02.05.94)</td><td>US</td></tr><tr><td>08/236,537</td><td>2 May 1994 (02.05.94)</td><td>US</td></tr><tr><td>08/412,383</td><td>28 March 1995 (28.03.95)</td><td>US</td></tr></table> (71) Applicant: THE ROCKEFELLER UNIVERSITY [US/US]; 1230 York Avenue, New York, NY 10021 (US). (72) Inventors: FISCHETTI, Vincent, A.; 448 Joan Court, West Hempstead, NY 11552 (US). CHEUNG, Ambrose; 500 East 63rd Street, New York, NY 10021 (US). (74) Agents: CRANE-FEURY, Sharon, E. et al.; Klauber & Jackson, 411 Hackensack Avenue, Hackensack, NJ 07601 (US).	08/227,514	14 April 1994 (14.04.94)	US	08/227,516	14 April 1994 (14.04.94)	US	08/236,536	2 May 1994 (02.05.94)	US	08/236,537	2 May 1994 (02.05.94)	US	08/412,383	28 March 1995 (28.03.95)	US	(81) Designated States: JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
08/227,514	14 April 1994 (14.04.94)	US														
08/227,516	14 April 1994 (14.04.94)	US														
08/236,536	2 May 1994 (02.05.94)	US														
08/236,537	2 May 1994 (02.05.94)	US														
08/412,383	28 March 1995 (28.03.95)	US														
(54) Title: PROCESS, APPARATUS AND REAGENTS FOR ISOLATING CELLULAR COMPONENTS																
(57) Abstract A reagent, method and container for the isolation of cellular components such as ribonucleic acid (RNA) from cells in a liquid solution. The container includes a cover assembly and a holder which is normally closed by the cover assembly and contains an RNA extractant solvent, micron-sized particles and at least one larger particle suitably of millimeter size. The container contains the reagent, which is an extractant solvent which contains phenol and guanidinium thiocyanate or guanidinium chloride and has a pH of about 4. The container also includes a friable sealing layer which separates the extractant solvent from the liquid medium containing the cells until the container is reciprocally shaken. The method includes the reciprocal shaking of the container, wherein the larger particle breaks the friable layer to permit mixing of the liquid medium with the extractant solvent resulting in the breaking of the cell walls by the micron-sized beads and the release of the RNA.																

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

1
PROCESS, APPARATUS AND REAGENTS FOR
ISOLATING CELLULAR COMPONENTS

BACKGROUND OF THE INVENTION

Field of the Invention

5 This invention relates to reagents, methods and apparatus for the isolation of cellular components such as ribonucleic acid (RNA) from natural cellular sources.

Description of the Related Art

Cells contain a wide variety of cellular components
10 appropriate to their function. They contain, for example, DNA, ribonucleic acid (RNA) and their expression products including a host of proteinaceous materials. This invention is useful for the isolation of such cellular components, but for convenience, the invention
15 will be described principally as it applies to the isolation of RNA.

RNA is a critical component in the sequence of biological reactions which results in the expression of the myriads of proteins including hormones, enzymes and structural
20 tissue essential for the existence of all forms of life. There is a critical need for large amounts of RNA for research purposes as well as diagnostic and therapeutic uses.

Plant cells and microorganisms including parasites, yeast
25 and, bacteria, are potential sources of RNA, but because the cell walls of RNA sources are so strong, it is difficult, time consuming and may require expensive equipment to utilize them as RNA sources.

RNA isolation from bacteria is difficult because the cell
30 walls are not readily susceptible to lysis. Current

protocols for isolating RNA from bacteria frequently employ enzymes such as lysostaphin or lysozyme to lyse the bacterial cells followed by the addition of denaturing agents to inactivate the ribonucleases.

- 5 However, these methods have not proven to be useful for the isolation of large amounts of high purity RNA.

U.S. Patent 4,843,155 employs an RNA isolation procedure in which biological tissue such as mouse anterior pituitaries are initially homogenized in a "glass-teflon
10 homogenizer" in an aqueous buffered medium at pH 4 containing phenol, a guanidinium salt such as guanidinium thiocyanate and, possibly an antioxidant. The homogenate is then extracted with an organic solvent insoluble in the buffer. Chloroform is exemplified as a useful
15 solvent. The RNA remains dissolved in the aqueous buffer. The DNA and protein of the biological tissue concentrate at the interface between the organic and inorganic phases. The RNA is separated from the aqueous phase by precipitation with a water soluble alkanol such
20 as isopropanol. The RNA may be recovered by centrifugation and removing of the supernatant. The patentee states that the isolation procedure can be completed in three hours.

This procedure is inconvenient because guanidine salts
25 are expensive and malodorous. Moreover, as described, the method is limited to biological tissue (i.e., eukaryotic and not prokaryotic cells) as an RNA source and requires an extended period of time to complete. The cells in biological tissue, being eukaryotic, are
30 enclosed by a fragile membrane which is easily ruptured.

Moreover, although glass beads have been used in methods of RNA isolation, glass beads cannot be used in conjunction with a homogenizer-type apparatus to achieve the results of the present invention, because beads would

cause the homogenizer to jam. Glass beads have been used in conjunction with a bead beater to isolate nucleic acids from cells. However, the disadvantage to this method lies in the fact that the method smashes the
5 cells, releasing the contents into medium in which unstable cellular components, such as RNA, are rapidly degraded by enzymes that are released.

Therefore, in view of the aforementioned deficiencies attendant with prior art methods of isolating cellular
10 components such as RNA from cells, it should be apparent that there still exists a need in the art for reagents, methods and apparatus suitable for the efficient isolation of such components from both eukaryotic and prokaryotic cells.

15 SUMMARY AND OBJECTS OF THE INVENTION

Accordingly, a major object of the present invention is to provide a method of isolating a cellular component including mechanically releasing such component by the application of reciprocal mechanical energy to a
20 suspension of cells in a liquid medium in a container which also contains a plurality of micron-size particles thereby to fracture the cell walls and release the cell component into the liquid medium and isolating the selected cell component from the medium.

25 Another object of the present invention is to provide a method of isolating RNA from a cellular source containing it together with DNA and protein including the steps of:
a) mechanically releasing the RNA from a cellular source containing it by the application of reciprocal
30 energy in the presence of an extraction solvent containing a plurality of micron size particles and comprising, by weight based on the total weight, from about 40 to 60% phenol in an aqueous buffer having a pH

of from about 4 to 4.5 which is 2M in guanidinium thiocyanate or guanidinium chloride;

b) adding a water insoluble organic solvent to form a two phase mixture including the aqueous extraction
5 solvent phase and an organic phase wherein the RNA is dissolved in the aqueous phase and the DNA and protein are concentrated in the organic phase or at the interface between the phases; and

c) precipitating the RNA from the aqueous phase by
10 the addition of a water soluble lower alkanol.

Still another object of the present invention is to provide a reagent useful for extracting RNA from a cellular source containing it together with DNA and protein which includes a mixture containing, by weight
15 based on the total weight of the mixture, from about 40 to 60% phenol and from about 0.1 to 1% of a detergent in an aqueous buffer having a pH of from about 4 to 4.5.

Yet another object of the present invention to provide a method of isolating RNA from a cellular source containing
20 it together with DNA and protein which includes the steps of:

a) mechanically releasing the RNA from a cellular source containing it in the presence of an extraction solvent containing a plurality of micron size particles
25 and comprising by weight based on the total weight, from about 40 to 60% phenol and from about 0.1 to 1% detergent in an aqueous buffer having a pH of from about 4 to 4.5;

b) adding a water insoluble organic solvent to form a two phase mixture including the aqueous extraction
30 solvent phase and an organic phase wherein the RNA is dissolved in the aqueous phase, the DNA and protein are concentrated in the organic phase or at the interface between the phases; and

c) precipitating the RNA from the aqueous phase by
35 the addition of a water soluble lower alkanol.

Another object of the present invention is to provide a container for isolation of ribonucleic acid (RNA) from RNA containing cells in a liquid medium which includes:

- (i) a holder having therein, an RNA extractant solvent, a plurality of micron-size particles and at least one larger particle;
- (ii) a cover assembly removably fastened onto the holder and having a cavity therein which is of sufficient volume to contain the liquid medium and the extraction solvent; and
- (iii) a friable sealing layer adapted to be broken by the larger particle upon reciprocal motion of the container, the sealing layer being adapted to separate the extractant solvent from the liquid medium until the sealing layer is broken by the larger particle.

An additional object of the present invention is to provide a container adapted to be firmly held in a reciprocal shaker useful for the extractive rupture of cells containing RNA to permit the extraction and recovery of the RNA therefrom, the container including:

- (i) a holder containing an extraction solvent and a plurality of particles of a particle size suitable for fracturing the walls of the cells from which the RNA is to be extracted; and
- (ii) a cover assembly for closing the container during reciprocal motion to prevent leakage of the contents from the container, the volume of the extraction solvent being sufficient to dissolve substantially all of the RNA released as a result of the fracture of the cell walls.

Also an object of the present invention is to provide a container adapted to be firmly held in a reciprocal shaker and useful for the extractive rupture of cells containing RNA to permit the recovery of RNA therefrom, wherein the container includes:

- (i) a holder containing an extraction solvent and a plurality of particles of a particle size suitable for fracturing the walls of the cells from which the RNA is to be extracted and at least one larger particle;
- 5 (ii) a cover assembly for closing the container thereby to prevent leakage of the contents from the container during reciprocal motion, the cover assembly having a cavity therein to contain the cellular source of RNA in a liquid medium; and
- 10 (iii) a friable seal positioned between the extraction solvent and the liquid medium, which seal is adapted to be broken during reciprocal motion by the impact of the larger particle thereby to permit mixing of the contents of the container.
- 15 Application serial number 08/227,514, entitled "Improved Method for Isolating RNA" describes and claims a method for extracting RNA from a natural source containing the RNA using reciprocal mechanical energy and micron size beads (particles) in an extraction solvent comprising
- 20 phenol in an aqueous buffer at pH of from 4 to 4.5, which is 2 molar in guanidinium thiocyanate or guanidinium chloride.

The invention described in Application Serial Number 08/227,516 entitled "Reagent and Process for Isolating

25 RNA" is similar to the novel procedure of the previous Application except for the use of a detergent in the buffered extraction solvent.

The process of this latter Application has all of the advantages of the process of Application Serial Number

30 08/227,514 and, in addition, utilizes an extractant which is much less expensive and more convenient to employ. Additionally, the process generally affords higher yields than the guanidinium process.

A common feature of both of the procedures of the two above-identified Patent Applications, in addition to the breaking of cell walls by mechanical energy from reciprocal motion, is the use of small solid micron-size beads (particles) to batter the cells and to break the cell walls like a bursting balloon. These particles are preferably glass beads, but may be any of a number of other particles such as metal, plastic or plastic coated metal particles.

10 The advantage of the process is that more cells are ruptured in a very short period of time so that more RNA is released to the extractant. In fact, the process can be completed in such a short period of time, just 30 seconds to 5 minutes, that no cooling is required, and no
15 special steps need be taken to be certain that the reagents employed are RNase free. RNase is a contaminating enzyme that causes the RNA to degrade.

The present invention likewise features a procedure which makes possible the separation and isolation of RNA of
20 exceptionally high purity in high yields. The procedure is very convenient and can be completed in a very short period of time, typically less than one half hour. The process is, moreover, applicable not only to biological tissue but also to microorganisms such as bacteria and
25 also to plant cells as sources of RNA. Such sources, especially bacteria are much more convenient than biological tissue as a source of RNA because they are uniform, readily available in any desired quantities and easier to work with than biological tissue.

30 In another aspect, this invention relates to novel containers for conducting processes for isolation of RNA or other cellular components. The containers comprise a cover assembly and a lower member for containing the

extractant and other components, hereinafter referred to as the "holder".

With the foregoing and other objects, advantages and features of the invention that will become hereinafter
5 apparent, the nature of the invention may be more clearly understood by reference to the following detailed description of the preferred embodiments of the invention and to the appended claims.

BRIEF DESCRIPTION OF THE DRAWINGS

10 Figure 1 is a side cross-sectional view of one embodiment of the present invention showing a container comprising a holder and its cover.

Figures 2-4 are side cross-sectional views of other embodiments of the containers of the present invention.

15 Figure 5 shows RNA purified from various mouse tissues using the FastPrep™ System, run on a native 1.2% agarose gel and stained with EtBr. Lane 1: liver; Lane 2: kidney; Lane 3: brain; Lane 4: spleen; Lane 5: muscle. Only the 18S and 28S rRNA bands are shown.

20 Figure 6 shows the results of RT-PCR on mouse liver RNA using B-actin primers. Samples were treated with DNaseI prior to amplification. Lane 1: 123 bp ladder; Lane 2: reverse transcription and amplification; Lane 3: amplification in the absence of reverse transcription.

25 Figure 7 shows the primer extension of the sarA transcript of *S. aureus*. RNA was purified using FastRNA™ reagent and FastPrep™ machine. The RNA was annealed to ³²P radiolabeled oligonucleotide primer and extended with reverse transcriptase, thereby yielding a primer
30 extension product. By running a sequencing reaction with

a DNA template (labeled GATC) and an identical primer, the transcription start site of the *sarA* transcript was mapped to position 101 in a *sarA* clone (see arrow). The samples 1 and 2 represent two different RNA preparations with the same method.

Figure 8 is a Northern blot of α -hemolysin transcript of parental strain ISP479C and car mutant 1E3. The size of the transcript was ≈ 1.8 kb. The OD₆₅₀ of 1.1 and 1.7 correspond to late-exponential and post-exponential phases, respectively.

Figure 9 is a Northern blot of protein A transcript in parent ISP479C, car mutant 1E3 and transductant 1E3-2. The agr mutant of ISP479C, strain B2, is included as a positive control. The protein A transcript is ≈ 1.9 kb. The arrow points to the 23S ribosomal RNA band which serves as a RNA marker.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS OF THE INVENTION

More particularly, the present invention relates to the application of sufficient mechanical energy to the cell walls of the selected RNA source in a particular extractant to disrupt the cell walls and release the RNA. The disruptive extraction process is effected in the presence of a chaotropic agent, i.e., a substance capable of disorganizing or disrupting molecular structures and molecular reactions within the cell.

The following are definitions of terms used throughout the present Specification and Claims:

The term "by weight", unless otherwise specified means by weight based on total weight.

"Reciprocal motion," "reciprocal energy," "reciprocal mechanical energy," "reciprocal agitation," "reciprocal shaking," and "reciprocally moving" are used interchangeably. These terms describe the use of a shaking motion, as opposed to rotational motion, in the presence of rapidly vibrating inert particles which are capable of fracturing the cell walls of prokaryotic organisms. The particles are preferably contained in a closed container, in a liquid medium. When the particles are "reciprocally agitated" in the presence of cells, two, opposing, colliding particles compress the cells in the middle and cause the cell walls to burst and release the cell components including RNA into the medium.

The term "liquid medium" can refer to a number of different media, including growth media, buffers or other liquids in which the cells are suspended.

The term "natural sources" means a source providing a biological sample of cells, including plant cells and tissues, eukaryotic cells and tissues, and prokaryotic cells.

The disruptive extraction process of the present invention is effected in the presence of, by weight, from about 40 to 60% phenol, in an aqueous buffer at a pH of from about 4 to 4.5 containing, as a chaotropic agent, sufficient guanidinium compound such as guanidinium thiocyanate or guanidinium chloride to provide a composition which is about 2 to 5 molar in the selected compound. Alternatively, the extractant may contain, by weight, 0.1-1% of a suitable surfactant. By disrupting the cells directly into an extraction medium, unstable cellular components such as RNA are immediately separated from proteins such as RNases, which would otherwise degrade the RNA.

In the process of the invention, cell walls of the RNA source, preferably bacteria are disrupted to release the RNA. Disruption is effected in an extraction solvent. The resulting mixture is then extracted with a water insoluble organic solvent. Since RNA is insoluble in organic solvents it remains in the aqueous phase. The DNA, proteins and other organic impurities either dissolve in the organic phase or collect at the interface between the layers. The separated RNA remains dissolved in the aqueous phase from which it may be precipitated by the addition of a water soluble lower alkanol such as methanol, ethanol or isopropanol. The thus isolated RNA may be recovered by filtration, centrifugation or other convenient procedure. The RNA, as recovered, is of exceptionally high purity and substantially free of DNA. The process is especially advantageous because of the high yields which can be obtained compared to other known processes.

Typically, useful sources include *Staphylococcus aureus*, *Streptococcus pyogenes*, *Mycobacterium tuberculosis*, *Neisseria meningitidis*, *Saccharomyces cerevisiae* (yeast), and *Chlamydomonas reinhardtii* (algae).

If desired, the DNA can be recovered in high yield and purity by conventional means well known to the skilled artisan.

Particularly important for the present invention is the discovery of the method of cell disruption in which the cell walls are fractured by force created by the reciprocal motion of the living cells and the cell wall fracturing particles in a closed container suspended in a liquid medium.

A novel reagent has now been discovered which is useful as an extraction solvent in this invention and makes

possible the separation and isolation of RNA of exceptionally high purity in high yields by a very convenient procedure which can be completed in a very short period of time, typically less than one half hour.

- 5 The novel reagents of this invention are mixtures containing, by weight, from about 40 to 60% phenol and from about 0.1 to 1% surfactant and an aqueous buffer having a pH of from about 4 to 4.5. At this pH, phenol is insoluble in aqueous media. As a result, the mixture
10 is a two phase mixture.

The process of the invention is applicable not only to biological tissue but also to microorganisms such as bacteria and also to plant cells as sources of RNA. Such sources, especially bacteria are much more convenient
15 than biological tissue as a source of RNA because they are uniform, readily available in any desired quantities and easier to work with than biological tissue.

If desired, the DNA can be recovered in high yield and purity by conventional means well known to the skilled
20 artisan.

There are a number of advantages to the process of this invention especially when conducted for the isolation of RNA. These include:

1. Applicability to RNA sources such as bacterial
25 cells and plant cells which have heretofore been refractory to homogenization procedures with other phenol extractant media.
2. Recovery of RNA as a high yield product substantially uncontaminated by DNA.
- 30 3. Applicability to the production of both small and large quantities of RNA in batch or continuous processes.

4. Completion in a very short period of time which makes it possible to conduct the process without cooling without RNase free reagents and without contamination by DNA fragments which would be produced by extended cell
- 5 disruption procedures.
 5. No ultracentrifugation is required.
 6. The novel reagents of the invention are substantially odor free and readily available at commercially attractive prices.
- 10 The invention also relates to novel containers for conducting processes for isolation of RNA or other cellular components.

In the course of cellular growth, a large variety of different RNA is produced for conversion and expression
15 of various proteins such as hormones, enzymes and other proteins. Following the expression of a required amount of such proteins, the cells would limit continued protein production by rapidly degrading the RNA coding for such protein. Accordingly, a maturing cell, at different
20 stages of its growth or activity will contain differing amounts of the same RNA molecules or differing amounts of different RNA molecules.

It thus becomes important to extract the RNA from the living cell at different stages during the growth cycle
25 of the cell to isolate and identify RNA species that may be short lived.

Certain of the embodiments of this invention are specially useful for these purposes. Thus, the cover assembly is adaptable to contain the growth medium,
30 including all the necessary nutrients, for cell growth. Optionally, the cells can be grown separately and aliquots of the growth medium, including the cells, can be transferred to the cover assembly prior to rupture of

the seal to permit mixing of the growth medium and cells with the extractive liquid. Alternatively, the cells can be isolated from a separate growth medium and added to the cover assembly in a buffer solution, with or without
5 an additive to stop, or induce, the production of a selected protein. Optionally, the cells can be grown separately and aliquots of the growth medium, including growth cells, can be transferred to the cover assembly prior to rupture of the seal to effect mixing of the
10 growth medium and the extraction liquid.

One embodiment of the container is especially useful for following the growth of cells by the isolation of various types of RNA produced in the course of cell development. In that embodiment, the cells of interest are cultured in
15 a suitable medium in a cavity formed in the cover assembly or cultured separately and placed in the cavity. The cover assembly or the holder has a friable liquid-tight seal which separates the extractant and the liquid medium containing the cells. The holder contains, in
20 addition to the extractant, a member of micron or sub-micron size particles and at least one large particle. When it is desired to extract the RNA the cover assembly is used to close the holder and the container is placed in a shaker. When the shaker is turned on, the resulting
25 reciprocal motion of the large particle fractures the seal and allows the liquid medium and extractant to mix. The motion of the micron size particles bursts the cell walls and releases the cell components including RNA.

In another embodiment, the container comprises a holder
30 which contains the extractant liquid, micron-sized beads and at least one larger bead in the extractant liquid, a friable seal, and a removable cover assembly. The cover assembly is removed, the cells in a liquid medium are placed on the friable seal and the cover closed. The

container is reciprocally agitated, for example by shaking, so that the larger bead breaks the seal.

The container of the present invention allows for cell disruption by the application of mechanical energy to obtain highly purified RNA from natural sources containing it together with DNA and other contaminants, chiefly proteins. One important aspect of the mechanical energy involved in cell disruption in accordance with this invention is that it is reciprocally applied, preferably in the presence of micron or submicron size beads, or equivalent projectile like objects, so as to smash the cell walls in the manner of a bursting balloon. Rotational energy such as generated with a blender or other homogenizer is not useful because the cells simply rotate and do not collide efficiently with each other and with the beads to allow the cells to be crushed between the beads.

The extraction solvent used is the guanidinium salt containing liquid described above.

Any of a wide variety of well known buffers which will permit control of the pH within the selected range may be employed. Buffers based on sodium acetate or sodium citrate are presently preferred because they are readily available and provide excellent results. Other buffers known to the skilled artisan may be used.

The extraction reagent containing the guanidine type chaotropic agent may optionally contain an antioxidant to limit the spontaneous oxidation of phenol and increase the shelf life of the extraction reagent. Typically useful antioxidants include organic antioxidants such as 2-mercaptoethanol, dithiothreitol, 2-hydroxy quinoline and cysteine.

- The first step in the practice of this invention is to mechanically fracture the cell walls of the cellular RNA source such as bacteria, the eukaryotic cells of biological tissue or plant cells. This may be
- 5 accomplished with any of a variety of available instruments of which presently preferred are bead beaters such as the Mini Bead Beater available from BioSpec Products of Bartlesville, OK. Other instruments are available.
- 10 Bead beaters are preferred instruments for treating the RNA source. Typically, the size of the bead is from about 0.1 to 1.0 microns, and will vary with the size of the DNA carrier cells. Even more preferable is the use of the FastPrep™ machine made by Savant Instruments
- 15 (Farmingdale, NY) because of its fast speed, reducing the time needed for complete disruption.

- One of the most important features of the selected shaker is that it is capable of generating enough mechanical energy by reciprocal motion to break the cell walls of
- 20 bacterial cells and release the RNA. This may be accomplished by rapidly vibrating glass beads or other inert particles in the extraction reagent in the presence of the bacteria or other RNA source. Such beads are commercially available in a variety of sizes from several
- 25 sources. They can be shaken at selected speeds to provide sufficient mechanical energy to disrupt the cell walls.

- RNA release is effected with the bacterial cells or other RNA source suspended in the extraction solvent. The time
- 30 required depends principally upon the size of the source from which the RNA is being extracted since the cell size of various RNA sources such as bacteria, yeasts and plant cells varies appreciably. For small batches of bacteria or other source, 30 seconds may be adequate. For larger

batches two or three minutes may occasionally be required. Time is a critical factor because the half-life of some of the RNA species is very short (1-2 minutes).

- 5 At the end of the disruption step when most of the RNA has been released into the buffered solvent, the solvent is extracted with an organic solvent which is substantially water insoluble. The solvent is added at the end of the extraction procedure. Typically useful
10 solvents include substituted and unsubstituted lower hydrocarbons such as chloroform.

During the extraction process the DNA dissolves in the organic solvent and may be recovered by standard techniques known to the skilled artisan. The RNA remains
15 dissolved in the aqueous buffer and may be precipitated with isopropanol as described above and illustrated in the example.

The present invention is especially useful when practiced with the novel reagent referred to above. The reagent is
20 a two phase mixture containing selected amounts of phenol and at least one detergent in an aqueous buffer at a pH of about 4. This pH is selected because when the hydrogen ion concentration is at this level, RNA is soluble in water and DNA is insoluble. A preferred
25 reagent is the FastRNA™ reagent manufactured by Bio 101. The selected detergent may be any of a variety of conventional surfactants including anionic, cationic, non-ionic and amphoteric surfactants.

Typically useful anionic detergents include, for example,
30 sodium lauryl sulfate, sodium-n-decyl sulfate and triethanolamine dodecyl benzene sulfonate.

Cationic detergents useful in the practice of the invention include, by way of example, cetyl trimethyl ammonium bromide and other N-alkyl quaternary ammonium halides, as well as polyethoxylated quaternary ammonium
5 chloride.

Amongst the nonionic detergents, there are tallow fatty alcohol ethoxylates, ethoxylated tridecyl alcohol, ethoxylated tridecanol, nonyl phenol ethoxylate and octylphenoxy polyethoxy ethanol.

10 Amphoteric detergents include, for example cocoamidopropyl betaine, disodium tallowimino dipriopionate and cocoamido betaine.

The surfactant which is presently the most preferred is cetyl trimethyl ammonium bromide because it gives the
15 best yields of DNA free product.

All of these detergents, and many other equivalent surfactant compounds are readily available from commercial sources.

As with the previous procedure, as of a number of buffers
20 or optional antioxidants may be employed.

The process of the invention when conducted with the novel extraction solvent is substantially similar to the process described above with the guanidinium salt.

As shown in Figure 1, the novel container which is
25 especially useful when employed in accordance with the invention to isolate RNA or other cellular components comprises a holder 10 and a cover assembly 11, both preferably of glass or an inert plastic. The holder contains a liquid RNA extractant 11, such as those
30 described above.

The holder 10 has preferably a cylindrical side wall so that it is round in sections perpendicular to Figure 1 and, like all embodiments of this invention, is adapted to be firmly held in a reciprocal shaker. A large number
5 of micron-size particles, suitably beads 12 are within the liquid extractant. Preferably the beads are of glass or metal and preferably they are from about 0.1-1.0 mm in diameter.

Any convenient number or weight of such beads may be
10 employed.

For research purposes, which normally require only small amounts of RNA the container may be very small. Typically, the total volume is from about 1.5-2.5 ml. Larger containers may be employed to obtain greater
15 quantities of RNA.

The holder is sealed with a friable seal 14 which may be of the same material as the holder itself and typically, is positioned about one-half way up the holder wall. The seal 14 is liquid-proof and yet light enough so that it
20 may be broken by the weight of the particles reciprocally moving in the container or, preferably by a larger particle such as bead 13. The larger bead may be of any convenient size and weight to fracture the seal.

The cover assembly 11 has internal screw threads 15 which
25 mate with the external screw threads 16 of container 10.

In the second, third and fourth embodiments shown in Figures 2-4, the holder 10, the extractant liquid 11, small-size beads 12 and larger bead 13, about 3-5 mm in diameter, are the same as in the first embodiment, shown
30 in Figure 1. In addition, the process steps used to isolate the RNA from the extractant liquid will also be the same.

In the second embodiment shown in Figure 2, the cover assembly 11A acts to store and nurture the cells. The cover assembly 11A has a ring-like side wall body member 20A, a friable liquid tight layer 21A which acts as the
5 bottom wall of the cavity 30, the cells of interest in a suitable medium 22A and a removable top cover 23A. The container consists of the holder 10, with its beads and extractant liquid, and the cover assembly 11A which is screwed, or otherwise fastened, onto the container body.

- 10 In use, when the user desires to isolate RNA from the cells in medium 22A, the cover assembly 11A will be screwed to holder 10 and with top cover 23A in place the container vigorously shaken for about 30 seconds to 5 minutes causing the large bead 13 to break the friable
15 layer 21A and the culture medium 22A or other liquid medium containing the cells will mix with the extractant liquid 11 and the other components in the holder.

In the embodiment of Figure 3, the cover assembly 11A has the cells of interest in liquid medium 22B. The cover
20 assembly has a ring-like sidewall 20B, cells of interest in a medium 22B, a solid top removable lid 23B and a solid bottom wall 24B. The container 10 has a friable liquid tight layer 21B, as in the first embodiment.

- In operation, the user will unscrew cover assembly 11B
25 from container 10, unscrew the top lid 23B and pour the medium 22B into the container 10, or transfer it by pipette, where it will be held on top of the friable layer 21B. The cover assembly 11B is then replaced on the holder and the container shaken so that
30 the bead 13 breaks the friable layer 21B, all of the components mix and the cell walls ruptured by the action of the beads, thereby to release the RNA.

In the fourth embodiment, shown in Figure 4, the cover assembly 11C has a ring-like side wall 20C, cells of interest in a culture or other medium 22C, a solid bottom wall 24C, and a solid top lid 23C and a top rim 25 having internal screw threads 26. The holder 10 has a friable liquid-tight layer 21C.

In operation, the user will remove cover assembly 11C from holder 10, unscrew the top lid 23C, and screw the cover assembly onto holder 10 with the screw threads 26. The container will then be shaken so that the bead 13 breaks the friable layer 21C.

As indicated above, the total volume of the container should be sufficient so that when it is closed, it will hold the extraction solvent, the liquid medium and the other components under conditions so that the entire mixture can be conveniently and efficiently shaken. A general rule for this purpose is that the total volume of the closed tube is one third occupied by beads, one third reagent/cells and 1/3 air space.

For use in research or other laboratories where relatively small amounts of RNA is required, the containers can be packaged in kits containing one or a plurality of containers together with a container of organic solvents and other accoutrements appropriate to the practice of the process. The kits may include a selection of containers with small beads of different sizes to accommodate the varying sizes of the cells employed as the RNA source. Such containers are especially useful with shakers which can hold a plurality of container, even up to 20 or more. Such machines and containers are especially useful when it is desired to conduct a number of RNA isolations simultaneously or sequentially.

The containers of this invention have been described principally with reference to four specific embodiments and to the isolation of RNA using an extraction solvent containing either a detergent or guanidium salt as the
5 chaotropic agent used to disrupt the molecular structure and molecular interactions in the cell containing the RNA to be isolated. Other chaotropic agents can be employed.

It will be apparent to the skilled artisan that the concept of the containers can be practiced utilizing
10 other structures that are equivalent to the structures illustrated and described. For example, the friable sealing layer could be fixed at the top of the holder so that it breaks when the cover assembly is tightly screwed into place thereby eliminating the need for large beads.
15 The same result could be achieved by sealing a pointed member on the inside of the cover assembly dimensioned so that it will fracture the sealing layer when the cover assembly is screwed tightly into place.

Of course, the cover assembly can be fixed to the holder
20 by means which are equivalent to the threads shown, for example by a force fit.

Since rupture of the cell walls releases all of the cellular constituents, this invention can be used with or without chaotropic agents and extraction solvents such as
25 those described above to isolate other cellular components using known isolation procedures. For example, proteins may be isolated from a disrupted mixture containing an extraction solvent that comprises a neutral buffer and a cocktail of protease inhibitors.

30 As another example, the processes and containers of the invention may be used to efficiently and rapidly shred tissue such as skin, intestine, gastric, liver etc. into the component parts for the isolation of certain

components. Individual cellular components, e.g., enzymes, may then be isolated using standard chromatographic techniques. Similarly, structural components e.g., connective tissue, membranes, cell wall components, etc. may be separated by differential centrifugation techniques.

The following examples are given by way of illustration only and should not be considered limitations of this invention, many apparent variations of which are possible without departing from the spirit or scope thereof.

EXAMPLES 1-6

RNA and DNA were isolated from the following microorganisms using the protocol described below:

1. *Staphylococcus aureus*
2. *Streptococcus pyogenes*
3. *Mycobacterium tuberculosis*
4. *Neisseria meningitidis*
5. *Saccharomyces cerevisiae* (yeast)
6. *Chlamydomonas reinhardtii* (algae)

Steps:

1. Centrifuge 10 ml of cells to be processed ($\sim 1-5 \times 10^8$ per ml).

2. Aspirate and discard the culture supernatant.

3. Resuspend bacterial pellet in 1 ml of extraction reagent.

Prepare 5 ml of the extractions by adding:

Phenol	2 ml (40%)
10% CTAB*	0.5 ml (1%)
2 M NaAc pH 4	125 μ l (50mM)
1 M DTT	μ l (1mM)
water	2.37 ml

* cetyl trimethyl ammonium bromide

This results in a two-phase reagent which needs to be vortexed vigorously before pipetting 1 ml of the reagent into the tube.

4. Transfer the reagent mixture into sterile microfuge
- 5 tube containing ≈ 0.5 ml (1.65 gm) of 0.1 mm glass/
Zirconia beads (larger beads used for yeast and algae).
5. Shake at 6000 rpm for 20 seconds with reciprocal
motion. Pulse spin to settle the glass/Zirconia beads to
the bottom of the tube.
10. 6. Add 200 μ l of chloroform. Let sit for 2 minutes.
7. Vortex for 25 seconds.
8. Centrifuge for 12,000 g for 15 minutes at 4°C.
9. Aspirate the upper aqueous phase.
10. Add 0.5 ml of isopropanol. Incubate for 10 minutes.
- 15 11. Spin 12,000 g for 10 minutes at 4°C.
12. Wash pellet with 70% ethanol and spin for 5 minutes
at 5000 x g.
13. Discard the 70% ethanol wash and air dry for 5-10
minutes.
- 20 14. Resuspend RNA for storage in 100 μ l of either
diethyl pyrocarbonate treated water or 0.5% sodium
dodecyl sulfate.
15. To obtain DNA, the interphase containing the DNA is
reextracted twice with phenol/chloroform (50/50 v). The
25 upper aqueous phase is saved. One-tenth volume of sodium

acetate (3M pH 5.2) is added. The DNA is precipitated with 3 volumes of 100% ethanol, centrifuged, washed with 70% ethanol and recentrifuged. The pellet containing DNA is resuspended in Tris EDTA buffer and used for
5 restriction digestion or cloning.

EXAMPLE 7

Fresh chunks of mouse liver, kidney, brain, spleen and skeletal muscle (~50 mg each) are placed in 2 ml tubes containing chaotropic RNA stabilizing reagents and a
10 lysing matrix and immediately processed in the FastPrep™ System for 30-60 seconds (There is no need for pre-homogenization). Following centrifugation, the clear RNA containing supernatant is ethanol precipitated, or matrix purified and eluted in 100 μ l DEPC H₂O. 10-20 μ l are run
15 on a native agarose gel (Figure 5). Figure 6 shows an RT-PCR analysis of liver RNA.

TABLE 1

TYPICAL RNA YIELD FROM VARIOUS SOURCES			
	SAMPLE	AMOUNT	TOTAL RNA YIELD
5	Muscle Tissue	100 mg	100-150 μ g
	Liver Tissue	100 mg	Up to 800 μ g
	Fibroblasts	10^7 Cells	50-80 μ g
	Lymphocytes	10^7 Cells	70-100 μ g
	Young Plant Leaf	250 mg	50-200 μ g
10	Mature Plant Leaf	250 mg	25-150 μ g
	Bacterial Cells	10^9 Cells	50-100 μ g
	Yeast Cells	10^9 Cells	50-100 μ g

EXAMPLE 8

A method for isolating RNA from gram positive bacteria which is fast, simple and requires no expensive equipment utilized the FastRNA™ (BIO 101, Vista, CA) kit together
5 with the FastPrep™ machine (Savant). The method was used for *Staphylococcus aureus* and *Saccharomyces cerevisiae* from which RNA extraction had previously been known to be difficult.

For this procedure, bacteria were cultured in growth
10 media overnight, diluted 1:100 in 10 mls of fresh media, and grown to mid log, late log and post-exponential phases corresponding to OD_{650nm} of 0.7, 1.1 and 1.7, respectively. The bacteria were then pelleted (2,500 x g for 10 minutes at 4°C) and resuspended in 1 ml of
15 reagent. The cells were transferred to a 2 ml microfuge tube, to which 500 µl of Zirconia/Silica beads (0.1 mm size) were added.

The tube was shaken in Savant machine for 20 seconds at 6500 rpm then placed on ice. 200 µl of chloroform was
20 added to the tube. After vortexing for 15 seconds and incubating at RT for 2 minutes, the mixture was centrifuged at 12,000 x g for 15 minutes at 4°C. The RNA was harvested from the aqueous phase (top layer), precipitated with 0.5 ml of isopropanol and pelleted
25 (12,000 x g for 10 minutes at 4°C). The pellet was washed with 75% ethanol, air-dried and resuspended in 100 µl of 0.5% SDS in DEPC-treated water. The RNA concentration was determined spectrophotometrically.

Five µg of each sample was electrophoresed through a 1.5%
30 agarose-0.66M formaldehyde gel in MOPS running buffer (20mM MOPS, 10mM sodium acetate, 2mM EDTA, pH 7.0). With this method, we were able to obtain a high yield of RNA with good purity as evidenced by the A₂₆₀/A₂₈₀ ratio (Table

1). This method of extraction is also highly efficient as the RNA yield appeared to be proportional to the number of bacterial cells (Table 1). The integrity of RNA species was also confirmed in formaldehyde gels (Figure 7).

Primer extension was carried out with 30 μ g of RNA resuspended in DEPC-treated water. The RNA was precipitated in the presence of the primer which was end-labeled (5×10^4 cpm) and then prehybridized overnight at 30°C. The reaction mixture was then precipitated with 5 volumes of ethanol in the presence of 10% acetic acid, washed and dried. Reverse transcription was carried out, and the reaction incubated at 42°C for 90 minutes, heated at 65°C for 0 minutes and then incubated with 1 μ l of RNase A for 15 minutes at 55°C. The reaction was cleaned up with QIAquick nucleotide removal kit (Qiagen), ethanol precipitated, dried and resuspended in 10 μ l of TE and 10 μ l of Sequenase stop solution. The reaction was heated to 95°C for 5 minutes, chilled on ice and loaded on sequencing gel.

EXAMPLE 9

Staphylococcal RNA was extracted from a 10 ml culture grown to late log or postexponential phase in CYGP was pelleted, resuspended in 1 ml of FastRNA™ reagent (BIO 101) and 0.5 ml of 0.1 mm Zirconia/Silica beads and shaken at 6,000 rpm in a high speed reciprocating homogenizer (FastPrep™ machine, Savant Instruments) for 20 seconds. [This apparatus is the subject of co-pending U.S. Patent Application Serial No. 08/294,544.] After adding 200 μ l of chloroform, the mixture was centrifuged at 12,000 g for 10 minutes. The RNA in the upper aqueous layer was precipitated with isopropanol and resuspended in DEPC-treated water. The total cellular RNA yield was

≈0.5 - 1 mg with an A_{260}/A_{280} ratio of 2.0. The integrity of the RNA species was verified with a formaldehyde gel.

Five μ g of RNA from each bacterial strain was applied to a 1.5% agarose-0.66 M formaldehyde gel in MOPS running
5 buffer as described in Cheung et al ((1994) *J. Bacteriol.* 176:580-585). RNA was transferred to Nytran nylon membrane using the Turboblottter System (Schleicher & Schuell, Keene, NH). RNA on the membrane was cross-linked with UV light, hybridized with a 32 P labeled gel-
10 purified protein A or α -hemolysin probe in 50% formamide at 42°C overnight, washed and autoradiographed as described in Cheung et al (1994).

WHAT IS CLAIMED IS:

- 1 1. A method of isolating a cellular component which
2 comprises mechanically releasing the component by the
3 application of reciprocal mechanical energy to a
4 suspension of cells in a liquid medium in a container
5 which also contains a plurality of micron-size particles
6 thereby to fracture the cell walls and release the cell
7 component into the liquid medium and isolating the
8 selected cell component from the medium.
- 1 2. The method of Claim 1 wherein the liquid medium
2 additionally contains a chaotropic agent.
- 1 3. A method of isolating RNA from a cellular source
2 containing it together with DNA and protein which
3 comprises the steps of:
 - 4 a. mechanically releasing the RNA from a cellular
5 source containing it by the application of reciprocal
6 energy in the presence of an extraction solvent
7 containing a plurality of micron size particles and
8 comprising, by weight based on the total weight, from
9 about 40 to 60% phenol in an aqueous buffer having a pH
10 of from about 4 to 4.5 which is 2M in guanidinium
11 thiocyanate or guanidinium chloride;
 - 12 b. adding a water insoluble organic solvent to form
13 a two phase mixture comprising the aqueous extraction
14 solvent phase and an organic phase wherein the RNA is
15 dissolved in the aqueous phase and the DNA and protein
16 are concentrated in the organic phase or at the interface
17 between the phases; and
 - 18 c. precipitating the RNA from the aqueous phase by
19 the addition of a water soluble lower alkanol.
- 1 4. The method of Claim 3 including the further step of
2 recovering the RNA by separating it from the aqueous
3 phase from which it has been precipitated by filtration

4 or by centrifugation and removal of the supernatant
5 aqueous phase.

1 5. The method Claim 1 wherein the extraction is
2 conducted in the presence of detergent or mixtures
3 thereof.

1 6. A reagent useful for extracting RNA from a cellular
2 source containing it together with DNA and protein which
3 comprises a mixture containing, by weight based on the
4 total weight of the mixture, from about 40 to 60% phenol
5 and from about 0.1 to 1% of a detergent in an aqueous
6 buffer having a pH of from about 4 to 4.5.

1 7. The reagent of Claim 6 wherein the detergent is
2 cationic.

1 8. The reagent of Claim 6 wherein the detergent is
2 cetyl trimethyl ammonium bromide.

1 9. A method of isolating RNA from a cellular source
2 containing it together with DNA and protein which
3 comprises the steps of:

4 a) mechanically releasing the RNA from a cellular
5 source containing it in the presence of an extraction
6 solvent containing a plurality of micron size particles
7 and comprising by weight based on the total weight, from
8 about 40 to 60% phenol and from about 0.1 to 1% detergent
9 in an aqueous buffer having a pH of from about 4 to 4.5;

10 b) adding a water insoluble organic solvent to
11 form a two phase mixture comprising the aqueous
12 extraction solvent phase and an organic phase wherein the
13 RNA is dissolved in the aqueous phase, the DNA and
14 protein are concentrated in the organic phase or at the
15 interface between the phases; and

16 c) precipitating the RNA from the aqueous phase by
17 the addition of a water soluble lower alkanol.

1 10. The method of Claim 9 including the further step of
2 recovering the RNA by separating it from the aqueous
3 phase from which it has been precipitated by filtration
4 or by centrifugation and removal of the supernatant
5 aqueous phase.

1 11. The method of Claim 9 wherein the detergent is
2 cationic.

1 12. The method of Claim 11 wherein the detergent is
2 cetyl trimethyl ammonium bromide.

1 13. The method of Claim 9 wherein the extraction is
2 conducted in the presence of detergents or mixtures
3 thereof.

1 14. A container for isolation of ribonucleic acid (RNA)
2 from RNA containing cells in a liquid medium which
3 comprises:

4 (i) a holder having therein an RNA extractant
5 solvent, a plurality of micron-size particles and at
6 least one larger particle;

7 (ii) a cover assembly removably fastened onto
8 the holder and having a cavity therein which is of
9 sufficient volume to contain the liquid medium and the
10 extraction solvent; and

11 (iii) a friable sealing layer adapted to be
12 broken by the larger particle upon reciprocal motion of
13 the container, the sealing layer being adapted to
14 separate the extractant solvent from the liquid medium
15 until the sealing layer is broken by the larger particle.

1 15. A container as in Claim 14 wherein the micron size
2 particles are glass beads about 0.1-1.0 mm in diameter.

1 16. A container as in Claim 14 wherein the combined
2 volume of the cavity and the holder is from about 1.5 to
3 3 ml.

1 17. A container as in Claim 14 wherein the extractant
2 liquid comprises a mixture of phenol and a chaotropic
3 agent in an aqueous buffer at a pH of from 4 to 4.5.

1 18. A container as in Claim 17 wherein the chaotropic
2 agent is a detergent.

1 19. A container as in Claim 17 wherein the chaotropic
2 agent is guanidium thiocyanate or guanidinium chloride or
3 detergent.

1 20. A container as in Claim 14 wherein the sealing layer
2 is within the holder.

1 21. A container as in Claim 14 wherein the cavity in the
2 cover assembly has a bottom wall and the sealing layer
3 forms at least part of the bottom wall.

1 22. A container adapted to be firmly held in a
2 reciprocal shaker useful for the extractive rupture of
3 cells containing RNA to permit the extraction and
4 recovery of the RNA therefrom, said container comprising:
5 (i) a holder containing an extraction solvent
6 and a plurality of particles of a particle size suitable
7 for fracturing the walls of the cells from which the RNA
8 is to be extracted; and
9 (ii) a cover assembly for closing the container
10 during reciprocal motion to prevent leakage of the
11 contents from the container, the volume of the extraction
12 solvent being sufficient to dissolve substantially all of
13 the RNA released as a result of the fracture of the cell
14 walls.

- 1 23. A container as in Claim 22 wherein the extractant
2 liquid comprises a mixture of phenol and a chaotropic
3 agent in an aqueous buffer at a pH of from 4 to 4.5.
- 1 24. A container as in Claim 23 wherein the chaotropic
2 agent is a detergent.
- 1 25. A container as in Claim 23 wherein the chaotropic
2 agent is guanidinium thiocyanate or guanidinium chlorite.
- 1 26. A container as in Claim 22 wherein the solid
2 particles are glass beads about 0.1-1.0 mm in diameter.
- 1 27. A container as in Claim 22 wherein the volume of the
2 holder is from about 1.5 to 2.5 ml.
- 1 28. A container as in Claim 22 further including a
2 friable sealing layer within the holder and at least one
3 millimeter size particle in the extractant liquid.
- 1 29. A container as in Claim 22 wherein the cover
2 assembly includes a cavity having a bottom wall and a
3 friable sealing layer which forms at least part of the
4 bottom wall.
- 1 30. A container adapted to be firmly held in a
2 reciprocal shaker and useful for the extractive rupture
3 of cells containing RNA to permit the recovery of RNA
4 therefrom, said container comprising:
- 5 (i) a holder containing an extraction solvent
6 and a plurality of particles of a particle size suitable
7 for fracturing the walls of the cells from which the RNA
8 is to be extracted and at least one larger particle;
- 9 (ii) a cover assembly for closing the container
10 thereby to prevent leakage of the contents from the
11 container during reciprocal motion, said cover assembly

12 having a cavity therein to contain the cellular source of
13 RNA in a liquid medium; and

14 (iii) a friable seal positioned between the
15 extraction solvent and the liquid medium, which seal is
16 adapted to be broken during reciprocal motion by the
17 impact of said larger particle thereby to permit mixing
18 of the contents of the container.

1 31. A container as in Claim 30 wherein the particles are
2 glass beads about 0.1-1.0 mm in diameter.

1 32. A container as in Claim 30 wherein the volume of the
2 holder is from about 1.5-3.0 ml.

1 33. A container as in Claim 30 wherein the extractant
2 liquid comprises a mixture of phenol and a chaotropic
3 agent in an aqueous buffer at a pH of from 4 to 4.5.

1 34. A container as in Claim 33 wherein the chaotropic
2 agent is a detergent.

1 35. A kit as in Claim 30 wherein the sealing layer is
2 within the container.

1 36. A kit as in Claim 30 wherein the cover assembly
2 includes a cavity having a bottom wall and the sealing
3 layer forms at least part of the bottom wall.

1 37. A method of isolating DNA from a cellular source
2 containing it together with RNA and protein comprising
3 the steps of:
4 a) mechanically releasing the RNA from a cellular
5 source containing it by the application of reciprocal
6 energy in the presence of an extraction solvent
7 containing a plurality of micron size particles and
8 comprising, by weight based on the total weight, from
9 about 40 to 60% phenol in an aqueous buffer having a pH

10 of from about 4 to 4.5 which is 2M in guanidinium
11 thiocyanate or guanidinium chloride;
12 b) adding a water insoluble organic solvent to
13 form a two phase mixture comprising the aqueous
14 extraction solvent phase and an organic phase wherein the
15 RNA is dissolved in the aqueous phase and the DNA and
16 protein are concentrated in the organic phase or at the
17 interface between the phases;
18 c) extracting the interphase with phenol and
19 chloroform; and
20 d) precipitating the DNA with ethanol.

FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8





FIG. 9



INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/04586

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07H1/08 C12N1/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07H C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 295 613 (MOORE W.E.C. AND BLANKS J.A.) 20 October 1981 see column 2, line 42 - line 54 ---	1
X	METHODS IN ENZYMOLOGY - GUIDE TO MOLECULAR CLONING TECHNIQUES, vol. 152, 1987 SAN DIEGO (US), pages 33-48, BERGER S.L. AND KIMMEL A.R. 'Section II General Methods for Isolating and Characterizing Nucleic Acids' see page 36 - page 37 ---	6
A	EP,A,0 288 618 (GEN-PROBE INCORPORATED) 2 November 1988 see the whole document --- -/--	3,6,9,37

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

5

Date of the actual completion of the international search 31 August 1995	Date of mailing of the international search report 07.09.95
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+ 31-70) 340-3016	Authorized officer Day, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 95/04586

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 554 034 (CHOMCZYNSKI P.) 4 August 1993 see the whole document -----	3,6,9,37

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 95/04586

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4295613	20-10-81	NONE	
EP-A-288618	02-11-88	AU-B- 604684	03-01-91
		AU-A- 7040487	24-09-87
		CA-A- 1296604	03-03-92
		JP-B- 5028599	26-04-93
		JP-A- 62236478	16-10-87
		JP-A- 5068553	23-03-93
		US-A- 5374522	20-12-94
EP-A-554034	04-08-93	US-A- 5346994	13-09-94
		JP-A- 5344886	27-12-93